

Information Sheet on Ramsar Wetlands

Categories approved by Recommendation 4.7 of the Conference of the Contracting Parties.

NOTE: It is important that you read the accompanying *Explanatory Note and Guidelines* document before completing this form.

1. Date this sheet was completed/updated:

26.10.98

For office use only.

dd mm yy

14/12/76 7IT014

Designation date

Site Reference Number

2. Country:

Italy

3. Name of wetland:

Sabaudia

4. Geographical coordinates:

41° 16' North; 13° 01' East

5. Altitude: (average and/or max. & min.). 0 m a.s.l.

6. Area: (in hectares) 1,474 ha

7. Overview: (general summary, in two or three sentences, of the wetland's principal characteristics)

It is a coastal lake with brackish water, adjoined on the seaward side by a sandy beach and recent coastal dunes. The lake itself is surrounded by limno-palustric deposits with peat soil, followed by inland dunes. The shore to the west is straight whilst that on the eastern side is interrupted by inlets penetrating the inland dunes.

8. Wetland Type: (please circle the applicable codes for wetland types as listed in Annex I of the *Explanatory Note and Guidelines* document)

marine-coastal: A - B - C - D - E - F - G - H - I - J - K

inland: L - M - N - O - P - Q - R - Sp - Ss - Tp
Ts - U - Va - Vt - W - Xf - Xp - Y - Zg - Zk

man-made: 1 - 2 - 3 - 4 - 5 - 6 - 7 - 8 - 9

Please now rank these wetland types by listing them from the most to the least dominant: J/Q, E, 4, Ts, 9

9. Ramsar Criteria: (please circle the applicable criteria; see point 12, next page.)

1a - lb - lc - 1d / 2a - 2b - 2c - 2d / 3a - 3b - 3c / 4a - 4b

Please specify the most significant criterion applicable to the site:

10. Map of site included? Please tick *yes* or *no*

(Please refer to the *Explanatory Note and Guidelines* document for information regarding desirable map traits)

11. Name and address of the compiler of this form:

Dr. S. Zerunian and Drssa. A. Noal, Ufficio Gestione ex ASFD di Sabaudia P. N. Circeo, Via Carlo Alberto 107, 04016 Sabaudia (LT). Tel. 0773-511385; Fax. 0773-510503.

Please provide additional information on each of the following categories by attaching extra pages (please limit extra pages to no more than 10):

12. Justification of the criteria selected under point 9, on previous page. (Please refer to Annex II in the *Explanatory Note and Guidelines* document)

Refer to attached bibliography.

13. General location: (include the nearest large town and its administrative region)

Coastal plain in the Lazio Region along the Tyrrhenian Sea and within the Province of Latina. The nearest towns are Latina (population 110,000) and Sabaudia (population 15,000).

14. Physical features: (e.g. geology, geomorphology; origins - natural or artificial; hydrology; soil type; water quality; water depth; water permanence; fluctuation in water level; tidal variations; catchment area; downstream area; climate)

Geology: The entire coastline is formed by a succession of clayey and clayey-sandy deposits with interbedding of peat soil and marshy deposits from the Quaternary era. The area is globally referred to by the term “red dunes” or old dunes, delimited to the west by a strip of recent dunes.

Geomorphology: The current situation has been influenced enormously by the major modifications which were performed during land reclamation works in the 1930’s. All the surface water has been collected in a network of channels by rectifying the routes of the water courses, pumping and filling depressions. This caused a complete transformation of the area, which was previously characterised by troughs and depressions (with differences in level of about 30m) due to the contraposition of old lines of dunes. At present, the coastal strip is relatively flat with man-made embankments which define both the water courses and the lakes.

Hydrology: The lake waters are brackish, tending towards salty, and the level of salinity varies between the summer and winter months. The channels collect the surface waters, with ducite characteristics. Some of the channels have an invert level lower than that of the sea, which causes the ascent of sea water for quite a long distance and a partial interference with the more superficial aquifer. Contamination with brackish water can therefore occur.

Land Types: The land types can be defined as follows:

- a) Non-agricultural land on recent dunes: Sand deposits which are strongly affected by the actions of the wind and salt. This gives rise to soils which are not very “mature”, lacking in organic substances and generally devoid of vegetation.
- b) Non-agricultural land on recent dunes: Similar to the previous land type but with a slightly greater content of organic substances. However, the land is still not very “mature”.
- c) Agricultural and pasture land: Fine lacustrine deposits mixed with fill material taken from the coastal dunes, giving rise to acidic peaty soils.
- d) Agricultural land: Reddish-yellow soils. Sandy, subacid soils resulting from Quaternary dunes, with little organic substance and clear signs of leaching.

Water Quality: The water table lies at a shallow depth, appearing at the surface at some points. The surface waters are mostly chlorides and alkaline-earth sulphates, and occasionally bicarbonate alkaline-earth. In some cases there is salt water contamination. It should be noted that the surface waters are subject to sudden changes in the values of the various parameters.

Fluctuations in Water Level: During the period immediately following the dry summer months there is a slight lowering of the water table due to low rainfall coinciding with intense exploitation of the water for irrigation purposes in the neighbouring areas. The fluctuation follows a cyclical trend that appears to be in equilibrium. However, what is more noticeable is the difference between the current level of the water table and that of 15 to 20 years ago, which shows a general lowering of the piezometric level as a result of the reclamation for agricultural use and the increase in water consumption.

Tides: The tide changes approximately every 6-6½ hours, with a range of between 12cm and 14cm.

Watershed: The numerous studies carried out have shown that the area is formed of a layer of sandy-clayey lenticular sediments which run parallel to the coastline. The presence of the interface fresh water-sea water makes it difficult to prepare a hydrological balance.

Downstream: Tyrrhenian Sea.

Upstream: Lepini, Ausoni and Aurunci mountain ranges. Water table lying on fractured calcareous material, with high secondary permeability which supplies a series of springs at the foot of the mountain ranges.

Climate: Mild, Mediterranean type climate, with rainfall concentrated in the autumn/winter period and relatively high temperatures. The summer months are dry. The rainfall between October and February generally corresponds to approximately 60% of the annual total whilst the rainfall between June and August reaches about 10%.

The average annual rainfall is approximately 900 mm. The driest month is July. The difference between the minimum and maximum monthly temperatures is quite low (usually less than 10°C). The humidity of the area is particularly high.

15. Hydrological values: (groundwater recharge, flood control, sediment trapping, shoreline stabilisation etc.)

See above.

16. Ecological features: (main habitats and vegetation types)

Behind the dunes: The vegetation is mostly arboreal with mixed, perennial, xeric species, including: *Ulmus minor*, *Populus sp.*, *Fraxinus exelsior*, *Juniperus oxicedrus*, *Pistacia lentiscus*, *Smilax aspera*, *Phyllirea latifolia*, etc.

Halophyte shores: These are occupied mostly by *Fragmitetia* and *Tifetia* (with partially flooded soil) situated near the fresh water pools and channels.

Fresh water marshes: In general, these areas were previously pasture lands and are characterised by the presence of species which adapt well to temporary flooding (after

rainfall). Herbaceous and grass species prevail, with clumps of *Juncus acutus* sometimes covering the entire surface area, which is also subject to periodic flooding.

Hedges and scrub: Rows of *Eucalyptus* and pine trees which were introduced during the land reclamation works.

17. Noteworthy flora: (indicating, e.g., which species/communities are unique, rare, endangered or biogeographically important, etc.)

The following species are found in the stagnant waters: *Phragmites australis*, *Calamagrostis pseudophragmites*, *Juncus effusus*, *J. acutus*, *J. inflexus*, *J. compressus*, *Scirpus maritimus*, *S. holoschoenus*, *S. lacustris*, *Cyperus longos*, *C. flavescens*, *Carex extensa*, *C. vulpina*, *Typha latifolia*.

The following species are found in the marshy areas with the greatest salinity: *Inula crithmoides*, *Atriplex hastata*, *Halimione portulacoides*, *Suaeda maritima*, *Limonium vulgare* and *Salicornia europaea*.

18. Noteworthy fauna: (indicating, e.g., which species are unique, rare, endangered, abundant or biogeographically important; include count data, etc.)

Aphanius fasciatus, *Knipowitschia panizzae* (*Padogobius panizzai*), *Rutilus rubilio*, *triturus carnifex*, *Rana italica*, *Bufo viridis*, *Hyla arborea*, *Testudo hermanni*, *Testudo graeca*, *Testudo marginata*, *Emys orbicularis*, *Podarcis muralis*, *Elaphe longissima*, *Coronella austriaca*, *Coluber viridiflavus*, *Natrix natrix*, *Natrix tessellata*.

In addition to the resident and breeding species there is a great deal of bird life, typical of lacustrine and marshy habitats, present during migration and in winter. The following breeding species are found: *Ixobrychus minutus*, *Gallinula chloropus*, *Cettia cetti*, *Cisticola juncidis*, *Acrocephalus scirpaceus*, *Acrocephalus arundinaceus*. The following species are found during migration and in winter: *Podiceps cristatus*, *Podiceps nigricollis*, *Tachybaptus ruficollis*, *Phalacrocorax carbo*, *Botaurus stellaris* (rarely), *Egretta garzetta*, *Ardeola ralloides*, *Ardea cinerea*, *Ardea purpurea*, *Nycticorax nycticorax*, *Plegadis falcinellus*, *Ciconia ciconia* (irregular during migration), *Ciconia nigra*, *Platalea leucorodia* (occasional), *Phoenicopterus ruber* (occasional), *Anser anser* (rarely), *Anas platyrhynchos*, *Anas penelope*, *Somateria mollissima* (rarely), *Mergus merganser*, *Mergus serrator*, *Fulica atra*, *Vanellus vanellus*, *Charadrius dubius*, *Charadrius hiaticula*, *Gallinago gallinago*, *Actitis hypoleucos*, *Tringa totanus*, *Tringa erythropus*, *Tringa nebularia*, *Calidris ferruginea*, *Larus argentatus*, *Larus melanocephalus*, *Larus ridibundus*, *Larus minutus*, *Chlidonias niger*, *Sterna sandvicensis*, *Sterna hirundo*, *Alcedo atthis*, *Cettia cetti*, *Gallinula chloropus*, *Acrocephalus melanopogon*, *Cisticola juncidis*, *Acrocephalus scirpaceus*, *Acrocephalus arundinaceus*, *Emberiza schoeniclus*.

Mammals: *Hystrix cristata*, *Vulpes vulpes*, *Meles meles*.

19. Social and cultural values: (e.g. fisheries production, forestry, religious importance, archaeological site etc.)

Social and recreational activities.

Fisheries.

Cheese products.

Archeological sites: numerous prehistoric finds have been made and there are significant, important Roman remains (villas, fountains, canals, statues).

20. Land tenure/ownership of: (a) site (b) surrounding area

(a) 70% Ministry of Agricultural Policies – ex ASFD management

30% Privately owned
(b) 100% Privately owned

21. Current land use: (a) site (b) surroundings/catchment

Extensive fishing industry.

Buffalo breeding.

Recreational, environmental education.

Bird-watching.

22. Factors (past, present or potential) adversely affecting the site's ecological character, including changes in land use and development projects: (a) at the site (b) around the site

The adverse factors have been greatly reduced following the inclusion of the area in the Circeo National Park. Previously, the zone was a hunting reserve and the lakes were exploited intensively for fishing and mussel farming. The greatest threat at present is the uncontrolled exploitation of the nearby areas for tourism which has caused a depletion of the plant cover of the coastal dunes (with consequent erosion phenomena) and the dumping of litter. The excessive human presence during the summer months has also caused pollution of the water table and the coastal waters.

23. Conservation measures taken: (national category and legal status of protected areas - including any boundary changes which have been made: management practices; whether an officially approved management plan exists and whether it has been implemented)

Inclusion in the Circeo National Park and successive expropriation order. The acquisition of the area has permitted a more precise conservation plan to be implemented, immediately eliminating the hunting (the area was previously a hunting reserve) and limiting human activities to ensure compatibility with conservation requirements.

Management plan: Islands are planned for bird life breeding, the restoration (at least partly) of the natural shoreline conditions, creation of small pools of freshwater (both permanent and temporary), control and revitalisation of the water system by dredging and possibly excavation of channels, pumping etc to avoid eutrophic phenomena.

Extensive fishing: Controlled fishing of bass, gilthead, eel, grey mullet, sole etc. is practised. Size of the catch is extremely variable depending on the season and the tides.

Animal breeding: One of the most widespread activities on the wetland pastures near the lakes is the breeding of buffalo, a species which has adapted perfectly well to the Pontine habitat. High quality dairy products are produced which are in great demand. With an extensive breeding it is possible to reconcile conservation and economic requirements.

Monitoring Units: Measurements are taken of temperature, humidity, wind speed and direction, rainfall, dissolved oxygen etc. enabling the state of "health" of the area to be evaluated.

24. Conservation measures proposed but not yet implemented: (e.g. management plan in preparation; officially proposed as a protected area etc.)

Improvement of water circulation in the lake.

Dredging of tidal canals.

Restoration of the banks and canals.

Formation of freshwater pools.

Preparation of nature trails and observation points on the banks.

25. Current scientific research and facilities: (e.g. details of current projects; existence of field station etc.)

Laboratory of nature biology and plant ecology.

Laboratory of hydrobiology.

Conference centre.

Exhibition hall for environmental subjects.

Accommodation for researches and students.

Library and cinema.

26. Current conservation education: (e.g. visitors centre, hides, information booklet, facilities for school visits etc.)

Botanical gardens.

Conference centre.

Exhibition hall.

Nature trails.

27. Current recreation and tourism: (state if wetland is used for recreation/tourism; indicate type and frequency/intensity)

Tourist activities are concentrated mostly in the spring and summer. The guided visits are organised by co-operatives, trails have been set-up (one of which has information boards for the blind) and signposts installed to increase awareness of the area.

28. Jurisdiction: (territorial e.g. state/region and functional e.g. Dept of Agriculture/Dept. of Environment etc.)

State: Italy; Region: Lazio; Province: Latina; Local Councils: Sabaudia and Latina.

29. Management authority: (name and address of local body directly responsible for managing the wetland)

Ministry for Agricultural, Food and Forestry Policies, management by ex ASFD., Via Carducci 5, Rome.

30. Bibliographical references: (scientific/technical only)

- Izzo G., Creo C., Grosso M., Silvestri C. *Indagine biogeochimica sui laghi costieri di Fogliano e Caprolace* (Biogeochemical studies on the coastal lakes of Fogliano and Caprolace).
- Cignini I., Falcucci M., Hull V., Sisi R. *Rappresentazioni modellistiche della dinamica trofica delle comunità planctoniche delle lagune del Parco nazionale del Circeo* (Dynamic trophic model of the plankton community in the lagoons of Circeo National Park).
- La Mura V., Spezie G. *Caratteristiche idrodinamiche della Laguna di Sabaudia. Risultati preliminari* (Hydrodynamic characteristics of the lagoons of Sabaudia – Preliminary results).
- Focardi S., Fossi M.C., Leonzio C., Lari L., Casini S., Corsolini S. *Applicazione di indicatori biologici della qualità delle acque nello studio del sistema dei laghi pontini* (Application of water quality biological indicators in a study of the Pontine lakes).
- Perdicaro R. *Livelli di metalli nella vongola verace del Lago di Fogliano* (Level of metals in the clams in Lake Fogliano).
- Mancini L., Dal Cero C., Volterra L. *Qualità microbiologica del lago di Fogliano ai fini del suo sfruttamento idrico* (Microbiological quality of Lake Fogliano relative to its use as a water resource).
- Fumanti B., Alfinito S., Cavacini P. *Osservazioni preliminari sul fitoplancton dei laghi di Fogliano, Monaci e Caprolace* (Preliminary observations on the phytoplankton of the lakes of Fogliano, Monaci and Caprolace).

- Giorgi U., Maccaroni A., Massa M., Ceccaroli C., Panella S. *Alcune considerazioni preliminari sulla comunità zooplanctonica dei Lago di Fogliano* (Preliminary observations on the plankton community of Lake Fogliano).
 - Zerunian S. *Indagine faunistica sui pesci del Parco nazionale del Circeo (Osteichthyes, Teleostei)* (Studies on the fish in Circeo National Park).
 - Gibertibi G., Zerunian S. *Studi sull'alimentazione del Cormorano nel P.N. del Circeo* (Study on cormorant feeding habits in Circeo National Park).
 - Massa F., Maccaroni A., Mariani A., Della Seta G., Giorgi U., Brizzi G., Panella S. *Scelte alimentari dei giovani di Mugilidi all'intero del lago di Fogliano* (Feeding of young Mugilidi in Lake Fogliano).
 - Minervini R. *Nuove forme produttive in vallicoltura: esperienze in corso presso l'Istituto Brunelli di Sabaudia* (New productive methods for lagoon fish breeding: experience of Brunelli Institute in Sabaudia).
 - Corbi F. *Risultati dei censimenti invernali degli uccelli acquatici nei laghi del P.N. del Circeo (1981-1995). Elementi per la gestione* (Results of census of wintering water-birds in the lakes of Circeo National Park 1981-1995).
 - Allavena S. *Le indicazioni della ricerca scientifica per la gestione del P.N. del Circeo* (Results of the scientific research for the management of Circeo National Park).
 - Anzalone B., Lattanzi E., Lucchese F., Padula M. *Flora vascolare del Parco nazionale del Circeo* (Vascular flora in Circeo National Park).
 - Biondi M., Pastorino A., Vigna Taglianti A. *L'avifauna nidificante del Parco nazionale dei Circeo* (Breeding birdlife in Circeo National Park).
-

Please return to: Ramsar Convention Bureau, Rue Mauverney 28, CH-1196 Gland, Switzerland
 Telephone: +41 22 999 0170 - Fax: +41 22 999 0169 - e-mail : ramsar @ hq.iucn.org